

Sani-Bowl

This product is a one-step disinfectant that is effective against a broad spectrum of bacteria, is virucidal* (including HIV-1 & HBV) and inhibits the growth of mold and mildew and their odors when used as directed.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Not for use on medical devices or medical equipment surfaces.

Disinfection, Virucidal* and Fungicidal Control

This is a proven one-step disinfectant, cleaner, sanitizer, fungicide, mildew stat, and virucide*.

Apply this product undiluted to walls, floors and other hard, (inanimate) non-porous surfaces such as tables, chairs, countertops, sinks, glazed tile, glazed porcelain and bed frames with a cloth, mop or mechanical spray device so as to thoroughly wet surfaces. Treated surfaces must remain wet for 10 minutes. If mechanical spray device is used, spray should be coarse. Spray 6-8 inches from surface. Rub with brush, sponge or cloth. Do not breathe spray (mist). Rinse with potable water after use on surfaces that come in contact with food.

NOTE: With spray applications cover or remove all food products. Prepare a fresh solution at least daily or when visibly soiled or diluted. For heavily soiled areas, a preliminary cleaning is required.

For heavily soiled areas, a preliminary cleaning is required. To disinfect hard, non-porous surfaces, treated surfaces must remain wet for 10 minutes. Then wipe excess with sponge, mop or cloth or allow to air dry.

Fungicidal against *Candida albicans* and *Trichophyton mentagrophytes* (the agent that causes athlete's foot fungus) on inanimate surfaces in locker rooms, dressing rooms and shower and bath areas and exercise facilities. Apply this product undiluted and follow the directions under "Disinfection, Virucidal* and Fungicidal Control".

Virucidal* Performance: This product was evaluated and found to be effective against the following viruses on inanimate hard non-porous environmental surfaces. Treated surfaces must remain wet for 10 minutes. For heavily soiled areas, a preliminary cleaning is required. Herpes Simplex Type1, Herpes Simplex Type2, HIV-1 (AIDS virus), Hepatitis B virus, Hepatitis C virus, Influenza A/Brazil Virus, Vaccinia virus.

Hepatitis B Virus: This product was proven to be effective against (Duck) Hepatitis B Virus with a contact time of 10 minutes.

HIV (AIDS virus): With a 2 minute contact time, this product was found to be effective against HIV-1 (AIDS virus).

Hepatitis C Virus: With a 10 minute contact time, this product was found to be effective against Bovine viral diarrhea virus (BVDV) (Surrogate for human Hepatitis C virus).

***KILLS HIV, HCV AND HBV ON PRECLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care setting or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of human immunodeficiency virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B virus (HBV) and Hepatitis C virus (HCV).

"SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HCV AND HBV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS."

PERSONAL PROTECTION: Clean up should always be done wearing protective latex gloves, gowns, masks and eye protection. (Specific barrier protection items to be used when handling items soiled with blood or body fluids are disposable latex gloves, gowns, masks, or eye coverings).

CLEANING PROCEDURE: Blood and other body fluids (containing HIV-1, HBV & HCV) must be thoroughly cleaned from surfaces and objects before application of this product.

DISPOSAL OF INFECTIOUS MATERIALS: Blood, body fluids, cleaning materials and clothing should be autoclaved and disposed of according to federal, state and local regulations for infectious waste disposal.

CONTACT TIME: For HIV, leave surfaces wet for two minutes. This contact time will not control other common types of viruses and bacteria. For HBV, leave surfaces wet for 10 minutes. For HCV, leave surfaces wet for 10 minutes.

3595 Zip Industrial Blvd
Atlanta, GA 30354
404-761-0657 Fax: 404-761-0709

Sani-Bowl

Fungicidal Activity: This product is effective against Trichophyton mentagrophytes (athlete's foot fungus) (a cause of Ringworm) and Candida albicans on inanimate, hard, non-porous surfaces in locker rooms, dressing rooms and shower and bath areas and exercise facilities. Apply this product undiluted and follow the directions under "Disinfection, Virucidal and Fungicidal Control".

Mold and Mildew Control Directions: This product will effectively inhibit the growth of mold and mildew (*Aspergillus niger*) and their odors on hard, non-porous surfaces. Thoroughly wet all treated surfaces completely. Let air-dry. Repeat application weekly or when growth or odor reappears.

NON-ACID TOILET BOWL (AND URINAL) DISINFECTION/CLEANER DIRECTIONS

Remove gross filth prior to disinfection.

Remove water from the toilet bowl. Apply 1-2 ounces of undiluted solution to the toilet bowl. Brush thoroughly over exposed surfaces and under the rim with a cloth, mop, or sponge. Allow to stand for 10 minutes and flush.

For Heavy Duty Cleaning: Empty toilet bowl or urinal and apply 8 ounces of product to exposed surfaces including under the rim with a cloth, mop, sponge or sprayer. Brush or swab thoroughly and allow to stand for 10 minutes and flush. For sprayer application, use a coarse spray device.

Cleaning/Deodorizing Directions: Apply undiluted solution with a mop, cloth, sponge or mechanical sprayer so as to wet all surfaces thoroughly. If mechanical spray device is used, spray should be coarse. Allow to air dry or wipe up excess liquid.

First Aid

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Causes moderate eye irritation. Harmful if swallowed or absorbed through skin. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking or using tobacco.

PHYSICAL OR CHEMICAL HAZARDS

Do not mix with oxidizers, anionic soaps and detergents.

STORAGE AND DISPOSAL

Store in original container in areas inaccessible to small children. Do not store on side. Avoid creasing or impacting of sidewalls. Do not reuse empty container. Wrap and discard in trash (or recycle).

3595 Zip Industrial Blvd
Atlanta, GA 30354
404-761-0657 Fax: 404-761-0709